

2010 AP Calculus AB - calculator

AB/BC 1)  $f(t) = 7t e^{\cos t}$  (rate snow accumulates from 12-9AM)

$g(t) = \begin{cases} 0 & 0 \leq t < 6 \\ 125 & 6 \leq t < 7 \\ 108 & 7 \leq t \leq 9 \end{cases}$  rates snow removed

a)  $\int_0^6 f(t) dt = \int_0^6 7t e^{\cos t} dt = \boxed{142.215}$

b) rate of accumulated snow - rate of removing snow (at 8AM)  
 $f(8) - g(8) = 7te^{\cos t} - 108 =$

$48.417 - 108 = \boxed{-59.583 \text{ ft per hour}}$

c)  $h(t) = \begin{cases} 0 & 0 \leq t < 6 \\ \int_6^t 125 & 6 \leq t < 7 \\ \int_6^t 108 & 7 \leq t \leq 9 \end{cases} \Rightarrow \begin{cases} 0 & 0 \leq t < 6 \\ 125(t-6) & 6 \leq t < 7 \\ 125 + 108(t-7) & 7 \leq t \leq 9 \end{cases}$

d)  $\int_0^9 7te^{\cos t} dt - \left[ \int_6^7 125 dt + \int_7^9 108 dt \right]$   
 amount accumulated - amount removed  
 $367.335 - (125 + 216) = \boxed{26.335 \text{ cubic ft}}$

$\boxed{3.67}$

AB/BC 2) a)  $\frac{E(7) - E(5)}{7 - 5} = \frac{21 - 13}{7 - 5} = \frac{8}{2} = 4$  400 entries per hour

b)  $\frac{1}{8} \left[ \frac{1}{2}(2)(0+4) + \frac{1}{2}(3)(4+13) + \frac{1}{2}(2)(13+21) + \frac{1}{2}(1)(21+23) \right]$   
 $\frac{1}{8} \left[ 4 + \frac{51}{2} + 34 + 22 \right] = \frac{1}{8} (85.8) = 10.688$

$\frac{1}{8} \int_0^8 E(t)$  is the average number of entries (in hundreds) per hour from noon until 8 PM.

c)  $\int_8^{12} t^3 - 30t^2 + 298t - 976$  (# processed from 8 AM - midnight)  
 $= 16$  (hundred) 1600

2300 (total entries) - 1600 (processed) = 700 left  
 at 8 PM between 8-12 TO process

d) processed most quickly when  $P(t)$  is at its max  
 ( $P(t)$  changes from increasing to decreasing)

This will occur at  $t=8$  or  $t=12$  or a critical point of  $P$ .

$P'(t) = 3t^2 - 60t + 298 = 0 \rightarrow t = 9.184, 10.817$

$P(8) = 0$ ,  $P(9.184) = 5.087$ ,  $P(10.817) = 291$

$P(12) = 8$

$t=12$  or midnight is when entries are processed most quickly.

$$3) a) \int_0^3 r(t) dt = \frac{1}{2}(2)(1000+1200) + \frac{1}{2}(1)(1200+800)$$

$$\textcircled{1} = 2200 + 1000 = \boxed{3200 \text{ people}} \textcircled{1}$$

b) Since  $r(t)$  is above 800 between 2 and 3 hrs, the number of people in line is increasing  $\textcircled{1}$   
(must have reason)

c) Line is longest when # of people in line reaches its max. This happens when  $r(t)$  changes from  $+$  to  $-$  (compared to 800 per hr). This

occurs at  $\boxed{t=3}$   $\textcircled{1}$

$$\textcircled{1} 700 + \int_0^3 r(t) dt - 800(3) \textcircled{1}$$

$$700 + 3200 - 2400 = \boxed{1500 \text{ people}} \textcircled{1}$$

$$d) 700 + \underbrace{\int_0^t r(t) dt}_{\textcircled{1}} - 800t = 0 \textcircled{1}$$

2010 AP Calculus AB no calculator

4) a) Area  $R = \int_0^9 (6 - 2\sqrt{x}) dx = \int_0^9 6 - 2x^{1/2}$   
 $= 6x - \frac{4}{3}x^{3/2} \Big|_0^9 = (54 - 36) - (0 - 0) = \boxed{18}$

b) Volume  $R = \pi \int_0^9 [(7 - 2\sqrt{x})^2 - (7 - 6)^2] dx$

c)  $V = \int_0^6 (\frac{y^2}{4} - 0) (3(\frac{y^2}{4} - 0)) dy$   $y = 2\sqrt{x}$   
 $\frac{y}{2} = \sqrt{x}$   
 $x = \frac{y^2}{4}$   
 $= \boxed{3 \int_0^6 (\frac{y^2}{4})^2 dy}$   $3 \int_0^6 \frac{y^4}{16} dy = \frac{3}{16} \int_0^6 y^4 dy$

5) a)  $g(3) = g(0) + \int_0^3 g'(x) dx = 5 + \frac{1}{4}\pi(2)^2 + \frac{1}{2}(1)(3)$   
 $5 + \pi + \frac{3}{2} = \boxed{\frac{13}{2} + \pi}$   
 $g(-2) = g(0) - \int_{-2}^0 g'(x) dx = \boxed{5 - \pi}$

b)  $y = g(x)$  has a point of inflection at  $x = 0, 2$  and  $3$ . This is where  $g'(x)$  changes from increasing to decreasing or decreasing to increasing.

\* c)  $h(x) = g(x) - \frac{1}{2}x^2$

$h'(x) = g'(x) - x$   $g'(x) = x$

At  $x=3$   $h'(x) = 3(x-2) - x = 2x - 6$  neg  $[2, 3)$

$h'(x) = -2x + 9 - x = -3x + 9$  neg  $(3, 5]$

$g'(x) = 3(x-2)$

$g'(x) = -1 - 2(4-5)$

neither a max or min at  $x=3$  since  $g'(x)$  doesn't change signs.

for  $[-2, 2]$   $x^2 + y^2 = 4$   $y = g'(x) = \sqrt{4 - x^2} = x$   $4 - x^2 = x^2$   $x = \sqrt{2}$   
 $h''(x) = g''(x) - 1$   $h''(\sqrt{2}) = g''(\sqrt{2}) - 1 < 0$  since  $g''(\sqrt{2}) < 0$

$\boxed{1.75}$

because  $g'(x)$  is decreasing from  $x=0$  to  $x=2$   
 It has a max at  $x = \sqrt{2}$  because  $g'(\sqrt{2}) = 0$   $g''(\sqrt{2}) < 0$

6) a)  $\frac{dy}{dx} = xy^3$   $f(1) = 2$   $m = 8$   $\boxed{y - 2 = 8(x - 1)}$

b)  $f(1, 1)$   $y - 2 = 8(1.1 - 1)$   
 $y - 2 = 8.8 - 8 \rightarrow y - 2 = .8 \rightarrow \boxed{y = 2.8}$   
 $\frac{d^2y}{dx^2} = 104 > 0$   $f(x)$  is concave up so  
 tangent line lies below.  
 Approx is less than actual value

c)  $\frac{dy}{dx} = xy^3$

$\int \frac{1}{y^3} dy = \int x dx$   
 $\frac{y^{-2}}{-2} = \frac{x^2}{2} + C$

$f(1) = 2$   $\frac{1}{2y^2} = \frac{x^2}{2} + C$   
 $-\frac{1}{8} = \frac{1}{2} + C$

$-\frac{5}{8} = C$

$-\frac{5}{2y^2} = \frac{x^2}{2} - \frac{5}{8}$

$\frac{4}{y^2} = -4x^2 + 5$

$y^2 = \frac{4}{-4x^2 + 5}$

$\boxed{y = \sqrt{\frac{4}{-4x^2 + 5}}}$  since  $f(1) = 2$  only positive